

Polarization transfer in SIDIS for Λ production[†]

Yuri Arestov

*Institute for High Energy Physics
142284 Protvino, Moscow Region, Russia*

Abstract. Polarization transfer from the longitudinally polarized positron to the final-state quark is considered for the underlying QCD subprocesses giving rise to the reaction $e_{\uparrow}^+ + p \rightarrow e^+ + \Lambda_{\uparrow} + X$ at the HERMES energy.

Measuring the longitudinal Λ polarization in the semi-inclusive deeply inelastic scattering (SIDIS)

$$e_{\uparrow}^+ + p \rightarrow e^+ + \Lambda_{\uparrow} + X, \quad (1)$$

that is under study in the experiment HERMES at DESY, may give important information for the determination of the twist-2 polarized fragmentation function $G_1(z, \mu)$. The latter can be found through the relation

$$D_{LL} \sim F_q(x) \cdot \hat{\sigma} \cdot \hat{d}_{LL} \cdot G_1(z, \mu), \quad (2)$$

where $F_q(x)$ is the unpolarized parton (quark/gluon) distribution function in the proton and the polarization transfer D_{LL} is defined as

$$D_{LL} = \frac{\sigma_{++} + \sigma_{--} - \sigma_{+-} - \sigma_{-+}}{\sigma_{++} + \sigma_{--} + \sigma_{+-} + \sigma_{-+}} \equiv \frac{d\Delta\sigma}{d\sigma}, \quad (3)$$

and it can be measured in the experiment. The hatted entities in (2) relate to the underlying QCD subprocess. The subscripts in σ 's in (3) denote the positron and Λ helicities.

The published HERMES measurement in the reaction (1) is $D_{LL} = 0.11 \pm 0.17 \pm 0.03$ [1].

As is seen from (2), it is very instructive to study $\hat{d}_{LL}$ for the purpose of the data handling and analysis. The parameter can also be expressed as $\hat{d}_{LL} = P_{out}^z / P_{in}^z$

[†] Talk given at IX Workshop on High Energy Spin Physics, Dubna, Aug. 2001

where P 's are the longitudinal components of the initial and final polarization vectors.

The elementary subprocesses under study are the elastic quark probing (Fig. 1)

$$e_{\uparrow}^+ + q \rightarrow e^+ + q'_{\uparrow} \quad (\text{LO}) \quad (4)$$

and the diagrams with the gluon correction (Fig. 2)

$$e_{\uparrow}^+ + q \rightarrow e^+ + q'_{\uparrow} + G \quad (\text{NLO}). \quad (5)$$

Since we are interested mainly in the central quark production region, here we neglect the contribution of the process $e_{\uparrow}^+ + G \rightarrow e^+ + q_{\uparrow} + \bar{q}$ with the quark




FIGURE 1. Elastic quark probing by polarized lepton.




FIGURE 2. The gluon emission by the quark line.

pair production by γ^*G coupling. The spin transfer properties in the production of heavy QQ quarks are discussed in ref. [2].

The purpose of this study is to look for kinematic regions where the polarization transfer $\hat{d}_{LL}$ gets enhanced in the subprocesses of fig. 1 and fig. 2. Being translated onto the hadronic level, these kinematic regions, if any, can serve as guidelines for the experiment HERMES. The interval of DIS variables (Q^2 , y and other) was chosen to be typical for HERMES.

The calculations are based on the following expression for the scattered quark vector polarization

$$P_{out}^z = \frac{(1 + P_z^{in})(|M_{++}|^2 - |M_{-+}|^2) + (1 - P_z^{in})(|M_{+-}|^2 - |M_{--}|^2)}{(1 + P_z^{in})(|M_{++}|^2 + |M_{-+}|^2) + (1 - P_z^{in})(|M_{+-}|^2 + |M_{--}|^2)},$$

which can be derived from the final-state spin-density matrix.

The properties of the elastic subprocess (4) are plotted in fig. 3. Both the differential cross section $d\sigma$ and polarized differential cross section $d\Delta\sigma$ decrease with increasing Q^2 so that the polarization transfer $\hat{d}_{LL}$ increases rapidly. It achieves the value ~ 0.8 at the extreme value of $Q^2 \sim 4 \text{ GeV}^2$. For modest momentum transfers, say $Q^2 \sim 2 \text{ GeV}^2$, $\hat{d}_{LL}$ has the 'normal' values of ~ 0.2 .

Fig. 4 represents the parameter $\hat{d}_{LL}$ calculated for the reaction (5) at four different values of the angle ϕ between two characteristic planes of the reaction – leptonic plane and the plane formed by the virtual photon and the outgoing quark. As is known, the polarization properties of the reactions $2 \rightarrow 3$ depend essentially, in general, on the rotational angle ϕ . The distributions of $\hat{d}_{LL}$ in fig. 4 are shown for




FIGURE 3. Elastic $e^+q \rightarrow e^+q$ scattering. (Q^2, y) dependences of the generated populations and differential cross section $d^2\sigma$ (left column); $d^2\Delta\sigma$ and parameter $\hat{d}_{LL}$ (right column).




FIGURE 4. The polarization transfer in the process $e^+q \rightarrow e^+Gq$ in the rest system of the virtual photon and incoming quark. p_z^* is the scattered quark momentum (in GeV). ϕ is the angle between (e^+, e^+) and (γ^*, q') planes.

four rotations of the planes with an increment equal to 90° . The lowest values of $\hat{d}_{LL}$ are obtained at $\phi = 180^\circ$ when the normals to the said plains are anti-collinear. The more sizeable values of $\hat{d}_{LL}$ appear when the normals make right angles (90° or 270°) – up to ~ 0.4 . And the most favorite kinematic region is predicted for the collinear normals ($\phi = 0$) where the polarization transfer increases up to 0.6.

So coming down to the experiment (1), one can expect the sizeable longitudinal component of the Λ polarization induced by the polarized positron under the kinematic configurations in which the normals to the leptonic plane and the plane (γ^*, Λ) are close to be collinear. The numeric estimates made at the HERMES energy can certainly be affected by the higher-order corrections.

REFERENCES

1. A. Airapetian et al, *Phys. Rev. D* (in press); e-print hep-ex/9911017.
2. Yu. Arestov, Twist-2 polarized fragmentation function in the open charm production in DIS, *Proc. Spin'2000*, Oct. 2000, Osaka, Japan.